

Letter to the Editor

Neurogenic pulmonary edema – the takotsubo syndrome of the lungs

Josef Finsterer, MD, PhD¹, Fulvio A Scorza, MD²

¹*Klinik Landstrasse, Messerli Institute, Austria.*

²*Neurological Department, Universidade de Sao Paulo, Brazil.*

Corresponding author: Dr. Finsterer J, Postfach 20, 1180 Vienna, Austria

Article history

Received 11 Feb 2021

Received in revised form 21 Feb 2021

Accepted 24 Feb 2021

Available online 28 Feb 2021

Keywords: Lung Edema, Stress Reaction, Takotsubo, Epilepsy, Brain-Lung Disease

This article reviewed by Dr. Prateek, Dr. Ram. Edited by Dr. Pradeep J., Dr. S Gaur. Available online 28 Feb 2021.

IJMS, all rights reserved.

Letter to the Editor

With interest we read the article by Cho et al. about a 31 years old male who experienced an inhalation injury and CO-poisoning after a fire in his apartment [1]. During hyperbaric oxygen therapy (HBOT), applied for the CO-poisoning, he experienced a single generalised tonic-clonic seizure followed by lung edema [1]. The case report has a number of shortcomings and raises concerns.

We do not agree with the conclusion that lung edema resulted from HBOT [1]. Acute lung edema after a generalised seizure may be the direct consequence of the seizure, as has been previously reported [2]. Acute lung edema immediately after an acute, severe neurological injury, such as acute meningitis, subarachnoid bleeding, seizures, ischemic stroke, intracerebral bleeding, epidural bleeding, traumatic brain injury, multiple sclerosis exacerbation, or vertebral artery dissection, is called neurogenic pulmonary edema (NPE) [3].

An argument in favour of NPE as a manifestation of a stress-reaction to a serious cerebral event is the simultaneous occurrence of NPE and TTS in a patient with meningioma [4], with ischemic stroke and thrombolysis [5], in patients with brainstem strokes [6], in a patient with epidural hemorrhage [7], in patients with subarachnoid bleeding [8], and in patients with brainstem hemangioblastoma [9].

A further shortcoming of the study is that no MRI of the brain was carried out. CO-poisoning is well known to cause hyperintensities on diffusion weighted imaging (DWI). In a retrospective study of 387 patients experiencing CO-poisoning, 26.9% had DWI hyperintensities on cerebral MRI [10]. DWI lesions could easily explain why the patients developed a seizure. DWI lesions may not show up on cerebral CT scans why carrying out an MRI is obligatory. Since CO-poisoning goes along with cerebral lesions, they are more plausible to explain the development of seizures than excessive oxygen consumption.

The main therapeutic options for treating NPE are supportive measures and lowering of the intracranial pressure [11]. Thus we should be informed if the index case had developed any clinical features or features on imaging of increased intracranial pressure. In case the patient had experienced an increase in intracranial pressure we should be informed which kind of treatment was applied.

In a recent study it has been shown that endothelin-1 may be elevated in the cerebrospinal fluid (CSF) of patients with NPE due to infection with enterovirus-71 [12]. We should be informed if endothelin-1 was elevated in the CSF of the index case. Diagnosing NPE is still challenging but transpulmonary thermo dilution studies can be helpful to establish the diagnosis. We are interested to know if this diagnostic approach was applied and how NPE

was diagnosed in the index case. Missing in the report is a description of the EEG.

Overall, this interesting case report could be more meaningful if an MRI of the brain had been carried out in the acute stage of CO-poisoning, if NPE triggered by the tonic-clonic seizure was more extensively discussed, and if treatment of NPE was more extensively described.

DECLARATION OF COMPETING INTEREST

There are no conflicts of interest.

FUNDING SOURCE

No funding was received.

USEFUL INFO

Author contribution: JF: design, literature search, discussion, first draft, critical comments.

Informed consent: was obtained.

The study was approved by the institutional review board.

REFERENCES

1. Cho K, Minami T, Okuno Y, Kakuda Y, Tsutsumi T, Kogame T, Ohtsuru S, Sato N, Koike K. Convulsive seizure and pulmonary edema during hyperbaric oxygen therapy: A case report. *J Med Invest* 2018; 65: 286-288.
2. July M, Williamson JE, Lolo D. Neurogenic pulmonary edema following non-status epileptic seizure. *Am J Med* 2014; 127: e9-10.
3. Al-Dhahir MA, Sharma S. Pulmonary Edema, Neurogenic, 2018 Oct 30. *Stat Pearls* [Internet]. Treasure Island (FL): Stat Pearls Publishing; 2018 Jan. Available from <http://www.ncbi.nlm.nih.gov/books/NBK532984/>.

4. Goyal A, Bharadwaj S. Neurogenic pulmonary edema and stunned myocardium in a patient with meningioma: A Heart-Brain Cross Talk. *J Neurosurg Anesthesiol* 2018. Doi: 10.1097/ANA.0000000000000492.
5. Kitagawa T, Yamamoto J, Kureshima M, Maeda H, Nishizawa S. Takotsubo Cardiomyopathy and neurogenic pulmonary edema following fibrinolytic therapy for embolic stroke: A Case Report. *No Shinkei Geka* 2018; 46: 21-25.
6. Probasco J C, Chang T, Victor D, Nyquist P. Isolated pulmonary edema without myocardial stunning in brainstem strokes. *J Neurol Transl Neurosci* 2014; 2: 1040.
7. Nakamori Y, Miyazawa N, Yoshitani K, Yamamoto S. A pediatric case with takotsubo cardiomyopathy and neurogenic pulmonary edema due to an epidural hemorrhage. *J Neurosurg Anesthesiol* 2018; 30: 279-280.
8. Manto A, De Gennaro A, Manzo G, Serino A, Quaranta G, Cancellà C. Early endovascular treatment of aneurysmal subarachnoid hemorrhage complicated by neurogenic pulmonary edema and Takotsubo-like cardiomyopathy. *Neuroradiol J* 2014; 27: 356-60.
9. Gekka M, Yamaguchi S, Kazumata K, Kobayashi H, Motegi H, Terasaka S, Houkin K. Hemorrhagic onset of hemangioblastoma located in the dorsal medulla oblongata presenting with tako-tsubo cardiomyopathy and neurogenic pulmonary edema: a case report. *Case Rep Neurol* 2014; 6: 68-73.
10. Jeon SB, Sohn CH, Seo DW, Oh BJ, Lim KS, Kang DW, Kim WY. Acute brain lesions on magnetic resonance imaging and delayed neurological sequelae in carbon monoxide poisoning. *JAMA Neurol* 2018; 75: 436-443.
11. Sari MY, Yıldızdaş RD, Yükselmiş U, Horoz ÖÖ. Our patients followed up with a diagnosis of neurogenic pulmonary edema. *Turk Pediatri Ars* 2015; 50: 241-4.
12. Tu YF, Lin CH, Lee HT, Yan JJ, Sze CI, Chou YP, Ho CJ, Huang CC. Elevated cerebrospinal fluid endothelin 1 associated with neurogenic pulmonary edema in children with enterovirus 71 encephalitis. *Int J Infect Dis* 2015; 34: 105-11.